Glossary of Terms Used in Medicinal Chemistry
(IUPAC Recommendations 1998)

A to H

Contents

Active transport*

Active transport is the carriage of a solute across a biological membrane from low to high concentration that requires the expenditure of (metabolic) energy.

Address-message concept

Address-message concept refers to compounds in which part of the molecule is required for binding (address) and part for the biological action (message).

ADME

Abbreviation for **Absorption, Distribution, Metabolism, Excretion**. (See also **Pharmacokinetics; Drug disposition)**.

Affinity

Affinity is the tendency of a molecule to associate with another. The **affinity** of a **drug** is its ability to bind to its biological target (**receptor**, **enzyme**, transport system, etc.) For pharmacological **receptors** it can be thought of as the frequency with which the **drug**, when brought into the proximity of a **receptor** by diffusion, will reside at a position of minimum free energy within the force field of that **receptor**.

For an **agonist** (or for an **antagonist**) the numerical representation of **affinity** is the reciprocal of the equilibrium dissociation constant of the ligand-**receptor** complex denoted K_A, calculated as the rate constant for offset (k_{-1}) divided by the rate constant for onset (k_1).

Agonist***

Catabolism consists of reactions involving endogenous organic substrates to provide chemically available energy (e.g., ATP) and/or to generate metabolic intermediates used in subsequent anabolic reactions.

Catabolite

A **catabolite** is a naturally occurring **metabolite**.

Clone*

A **clone** is a population of genetically identical cells produced from a common ancestor. Sometimes, "**clone**" is also used for a number of recombinant DNA (deoxyribonucleic acid) molecules all carrying the same inserted sequence.

Codon*

A **codon** is the sequence of three consecutive **nucleotides** that occurs in mRNA which directs the incorporation of a specific amino acid into a protein or represents the starting or termination signals of protein synthesis.

Coenzyme

A **coenzyme** is a dissociable, low-molecular weight, non-proteinaceous organic compound (often **nucleotide**) participating in enzymatic reactions as acceptor or donor of chemical groups or electrons.

Combinatorial synthesis

Combinatorial synthesis is a process to prepare large sets of organic compounds by combining sets of building blocks.

Combinatorial library

A **combinatorial library** is a set of compounds prepared by combinatorial synthesis.

CoMFA

See **Comparative Molecular Field Analysis**.

Comparative Molecular Field Analysis (CoMFA)**

Comparative molecular field analysis (CoMFA) is a **3D-QSAR** method that uses statistical correlation techniques for the analysis of the quantitative relationship between the biological activity of a set of compounds with a specified alignment, and their three-dimensional electronic and steric properties. Other properties such as hydrophobicity and hydrogen bonding can also be incorporated into the analysis. (See also **Three-dimensional Quantitative Structure-Activity Relationship [3D-QSAR]**).

Computational chemistry**

Computational chemistry is a discipline using mathematical methods for the calculation of molecular